

TABLE II—U.S. GRADE B OR U.S. CHOICE: ALLOWANCES FOR DEFECTS—Continued

| Total allowance | | Limitations | |
|---|--|-------------|--|
| Dirt. Foreign material. Insect infestation. Decay. | Foreign material. Insect infestation. Decay. | | |

TABLE III—U.S. GRADE C OR U.S. STANDARD ALLOWANCES FOR DEFECTS

| Total allowance | | Limitations | |
|--|--|---|--|
| Not more than a total of 20 percent, by weight, may be damaged or affected by: Off-color. Poor texture. End cracks. ¹ Skin or flesh damage. ² Fermentation. Scars. Heat damage. Insect injury. Other means. Mold. Dirt. Foreign material. Insect infestation. Decay. | Not more than 10 percent, by weight, may be damaged or affected by: End cracks. ¹ Skin or flesh damage. ² Fermentation. Scars. Heat damage. Insect injury. Other means. Mold. Dirt. Foreign material. Insect infestation. Decay. | Not more than 8 percent, by weight, may be damaged or affected by: Skin or flesh damage. ² Fermentation. Scars. Heat damage. Insect injury. Other means. Mold. Dirt. Foreign material. Insect infestation. Decay. | Not more than 5 percent, by weight, may be affected by: Mold. Dirt. Foreign material. Insect infestation. Decay. <i>Provided, That not more than 1 percent, by weight, may be affected by decay.</i> |

¹ Except that each 1 percent of end cracks to, and including, 8 percent, by weight, shall be considered as ½ percent damaged by end cracks; and any additional end cracks shall be calculated as true percentage, by weight.

² Not applicable to "Whole Pitted" style.

TABLE IV—MOISTURE ALLOWANCES FOR DRIED PRUNES

[Non-hermetically sealed containers; 10 pounds or more]

| Grades | Maximum moisture limits (percent) | |
|-------------------------------------|---------------------------------------|---------------------------------------|
| | Counts averaging 60 or less per pound | Counts averaging 61 or more per pound |
| U.S. Grade A or U.S. fancy | 25 | 24 |
| U.S. Grade B or U.S. choice | 25 | 24 |
| U.S. Grade C or U.S. standard | 25 | 24 |
| Substandard | 25 | 24 |

[21 FR 8177, Oct. 25, 1956, as amended at 30 FR 11596, Sept. 10, 1965. Redesignated at 42 FR 32514, June 27, 1977, and further redesignated at 46 FR 63203, Dec. 31, 1981]

MOISTURE, UNIFORMITY OF SIZE, DEFECTS

§ 52.3185 Moisture limits.

Dried prunes shall not exceed the moisture limits for the applicable grades and kind and size of packaging as designated in Table IV of this subpart except there is no moisture limit when safe and suitable preservatives

have been added. "Moisture" means the percentage by weight of the finished dried prunes, exclusive of pits, that is moisture when determined by the Dried Fruit Moisture Tester Method or in accordance with methods that give equivalent results. The moisture limits in Table IV apply only to so-called "bulk packs" of dried prunes packaged in non-hermetically sealed containers holding 10 pounds or more of dried prunes when safe and suitable preservatives have not been added. Such containers include, but are not limited to, wood boxes or fiber boxes.

[30 FR 11596, Sept. 10, 1965. Redesignated at 42 FR 32514, June 27, 1977, and further redesignated at 46 FR 63203, Dec. 31, 1981]

§ 52.3186 Definitions for uniformity of size.

(a) *Fairly uniform in size.* "Fairly uniform in size" means that in a sample of 100 ounces:

(1) For prunes that average 53 prunes or less per pound, the count per pound of 10 ounces of the smallest prunes does not vary from the count per pound of 10